

Black Soldier Fly Larvae – A Sustainable Protein Source?


AUTHOR DETAILS: Ian Banks is currently studying a PhD in Entomology and Sanitation at the London School of Hygiene and Tropical Medicine. The project aims to evaluate the potential of black soldier fly (*Hermetia illucens*) larvae, as agents to digest pit latrine contents, and discuss the impact of this approach on sanitation in developing countries.

AQUACULTURE & FISHMEAL

Aquaculture

- Aquaculture is a growing source of food for the human race
- In 2008, aquaculture produced 43%, 52.5 million tonnes (Mt)¹, of the total fish globally caught and produced
- Estimates predict that by 2050, between 95 and 167 Mt of fish will need to be produced using aquaculture, depending on consumption, and assuming global capture remains at 2008 levels²

Predicted Aquaculture Production 2008 - 2050




- The production of aquaculture fish is highly dependent on a sufficient supply of nutrients, with up to 80% being fishmeal³

Fishmeal

- Fishmeal is a brown powder produced by cooking, pressing, and drying fish and fish trimmings

- Fishmeal contains high levels of protein (60 - 72%) and fat (5 - 12%), with a high content of health promoting omega-3 fatty acids⁴
- In 2008, 20.8 Mt of fish were caught and used to produce 4.8 Mt of fishmeal^{1,5}, with 58% used in aquaculture

Global Fishmeal Production & Use in Aquaculture 1988 - 2008



- The cost of fishmeal has increased dramatically since 2000, due to static global supplies and strong market demand³



Cost of Fishmeal 1982 - 2012

- Therefore, alternative sources of nutrients used in aquaculture will be required to replace, partially or fully, fishmeal

Fishmeal Alternative

- There is a large variety of alternative protein sources available that could be used to replace fishmeal, including; plant by-products, animal by-products, and invertebrates (Table 1)

Table 1 - Typical Composition of Fishmeal Replacements

Origin	Ingredient	Protein (%)	Lipid (%)	Lysine (%)	Methionine (%)	Cysteine (%)	Advantages	Disadvantages
Plant	Soybean meal (SBM) ⁶	48.5	0.9	3.08	0.68	0.75	Economical, nutritious, high crude protein	Concentrations of 10 essential amino acids (EAA), fat, ash and phosphorous are low
	Canola meal ⁶	38	3.8	2.27	0.7	0.47	Not widely used in aquafeed, similar protein content to SBM	Similar price to SBM, low in phosphorous
Animal	Poultry by-product meal ⁷	66	13	3.10	0.99	0.98	High crude protein and Lysine	Deficient in Lysine, Methionine and Histidine
	Fish by-product meal ⁸	Varies depending on species					Best nutritional substitute	Potential pathogens and contaminants harmful to both fish and humans
Invertebrate	Earthworm meal ⁹	63	18.5	1.43	1.72	0.39	High crude protein, and fat	Low Lysine and Cysteine
	Housefly pupae meal ¹⁰	41	8.5	6.04	2.28	0.52	High crude protein, Lysine and Methionine	Low Cysteine and fat, disease spreading vector,
	Black soldier fly meal ¹¹	52.5	19.2	2.16	0.32	0.31	High crude protein, fat and Lysine, can alter EAA concentrations with different diets	Low Methionine and Cysteine

AQUACULTURE & FISHMEAL

- The black soldier fly (BSF) (*Hermetia illucens*) is a non-disease spreading, non-pest fly that is found around the world¹²

Adult black soldier fly (*Hermetia illucens*)



- The first five larval instars are voracious consumers of organic matter, the sixth larval instar is known as the prepupae and does not feed¹³
- These prepupae are high in fat and protein, and when used as a component of a complete diet support good growth in chickens¹⁴, swine¹⁵, rainbow trout¹⁶, and catfish¹⁷

Black soldier fly prepupae



- It is possible to enhance BSF meal with Omega-3 fatty acids¹⁶, and essential amino acids¹¹, by altering what the black soldier fly larvae feed on, this meal can replace up to 50% of fishmeal without affecting fish growth¹¹
- It is estimated that with a feed conversion rate (FCR) of 16%, BSF feeding on swine manure in the United States could produce 1.8 Mt of prepupae per year
- Research is being conducted on the efficiency of BSF's converting fresh human faeces and pit latrine waste into prepupal biomass, with preliminary results producing an FCR of at least 16%¹⁸
- With 1.7 billion people using latrines, and a further 2.7 billion with no access¹⁹, it is estimated that 25 Mt of prepupae could be produced per year from the waste created

CONCLUSION

- Aquaculture is set to grow greatly over the next 40 years, while its main source of nutrition is unsustainable and increasing in price
- Black soldier fly larvae offer a suitable fishmeal alternative, while at the same time providing a way to effectively manage human and animal waste



References

- 1) Food and Agriculture Organization of the United Nations (FAO), (2010). The State of Food Insecurity in the World: Addressing food insecurity in protracted crisis.
- 2) Allen, G., (2008). Sustainable Aquaculture Feeds, Industry & Investments NSW. Port Stephens Fisheries Institute, pp. 3.
- 3) Tacon, A.G.J., & Metten, M., (2008). Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: Trends and future prospects. *Aquaculture*, **285**, pp 145 - 158.
- 4) International Fishmeal and Fish Oil Organization (IFFO), (2011). What are fishmeal and fish oil?
- 5) FAO FishStatJ, (2012). FishStatJ - Universal Software for Fishery Statistical Time Series.
- 6) Gatlin, D.M., et al., (2007). Expanding the utilization of sustainable plant products in aquafeeds: a review. *Aquaculture Research*, **38**, pp 551 - 579.
- 7) Laporte, J., et al., (2009). Poultry meat meal: a valuable source of the partial replacement of fish meal in aquafeeds. *International Aquafeed*, **12**, (2), pp 12 - 17.
- 8) Hardy, R., (2004). Problems and opportunities in fish feeds: fisheries processing byproducts. *International Aquafeed*, **7**, (2), pp 33 - 34.
- 9) Istiqomah, L., et al., (2009). Amino Acid Profile of Earthworm and Earthworm Meal (*Lumbricus rubellus*) for Animal Feedstuff. *J. Indonesian Trop. Anim. Agric*, **34**, (4), pp 253 - 257.
- 10) Aniebo, A.O., et al., (2009). Replacement of fish meal with maggot meal in African catfish (*Clarias gariepinus*) diets. *Revista Cientifica UDO Agrícola*, **9**, (3), pp 666 - 671.
- 11) Seeley, W.M., et al., (2011). Sensory Analysis of Rainbow Trout, *Oncorhynchus mykiss*, Fed Enriched Black Soldier Fly Prepupae, *Hermetia illucens* Journal of the World Aquaculture Society, **42**, (1), pp 34 - 45.
- 12) Loizelet, M. (1969). Dispersion et transport des insectes nuisibles: *Hermetia illucens* L. (Diptera: Stratiomyidae). *Bulletin des recherches agronomiques de Gembloux*, **4**, (1).
- 13) Copello, A., (1926). Biologia de *Hermetia illucens* Lat. *Journal of the Entomological Society Argentina*, **3**, pp 23 - 27.
- 14) Hale, O. M. (1973). Dried *Hermetia illucens* larvae (Diptera: Stratiomyidae) as a feed additive for poultry. *Journal of the Georgia Entomological Society*, **8**, pp 16 - 20.
- 15) Newton, G. L., et al., (1977). Dried *Hermetia illucens* Larvae Meal as a Supplement for Swine. *Journal of Animal Science*, **44**, (3), pp 395 - 400.
- 16) St-Hilaire, S., et al., (2007). Fly prepupae as a feedstuff for rainbow trout, *Oncorhynchus mykiss*. *Journal of the World Aquaculture Society*, **38**, (1), pp 59 - 67.
- 17) Newton, G. L. (2005a). The black soldier fly, *Hermetia illucens*, as a manure management/resource recovery tool. *Proceedings of the Symposium on the State of the Science of Animal Manure and Waste Management*, January 5-7, 2005, San Antonio, Texas, USA.
- 18) Banks, I.J., (2011). Unpublished Data. London School of Hygiene and Tropical Medicine.
- 19) WHO, UNICEF, et al., (2010). Progress on sanitation and drinking water - 2010 update. Geneva, New York.

BILL & MELINDA
GATES foundation

AgriProtein
technologies

Water
SHED
Water, Sanitation & Hygiene Enterprise Development